



Memovit B12 SYRUP

DL-PHOSPHOSERINE - CYANOCOBALAMINE - L-GLUTAMMINE

COMPOSITION

Each 10 g vial contains:

Active principles: DL-phosphoserin 40 mg, cyanocobalamin 500 mcg.

Excipients: sorbitol 70%, raspberry aroma, methyl P-oxybenzoate, distilled water.

Each proportioner stopper contains:

Active principle: L-glutamine 60 mg

Excipient: mannitol.

PHARMACEUTICAL FORM AND CONTENT

6 syrup vials with proportioner stopper

10 syrup vials with proportioner stopper

PHARMACEUTICAL CATEGORY

Memovit B12 represents an association of three substances known for their favourable action on nervous tissue metabolism acting in neurotronic and neurotrophic sense.

MARKETING AUTHORISATION HOLDER

MAGIS FARMACEUTICI S.r.l. – Via Cefalonia, 70 - 25124 Brescia - Italy

PRODUCER AND FINAL CONTROLLER

Doppel Farmaceutici S.r.l. - Via Martiri delle Foibe, 1 – 29016 Cortemaggiore (PC).

Sales concessionaire: AESCULAPIUS FARMACEUTICI S.r.l. – Brescia - Italy

THERAPEUTIC INDICATIONS

Intellectual overtiring states with lack of appetite, asthenia, easy exhaustability.

CONTRAINDICATIONS

Individual known hypersensitivity to the components.

SPECIAL WARNINGS

Though this preparation contains cyanocobalamin (B12 vit.) it cannot substitute the parenteral administration of cyanocobalamin in the treatment of pernicious anemia.

The product can cause slight excitability and insomnia; therefore it is not advisable the administration in the evening.

After a short period of treatment without appreciable results it's necessary to consult a doctor.

Memovit B12 contains sorbitol and parabens

This medicine contains 1,75 g of sorbitol per dose.



Sorbitol is a source of fructose. If your doctor has told you that you are intolerant to some sugars, or if you have a diagnosis of hereditary fructose intolerance, a rare genetic disorder where patients are unable to transform fructose, talk to your doctor before you take this medicine. .

If your doctor has diagnosed you have an intolerance to some sugars, contact him before taking this medicine.

In addition, the medicine contains parabens (preservatives) such as methyl para-hydroxybenzoate which can cause allergic reactions (even delayed).

DOSAGE AND METHOD OF ADMINISTRATION

One vial a day, preferably in the morning on an empty stomach. Take off the capsule pulling up the tongue and press on the jutting stopper till you obtain the powder to fall into the vial. Then agitate till you get a complete solution, pour in a glass and add water at pleasure.

POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Reporting of side effects

If you experience any side effects, including those not listed in this leaflet, talk to your doctor or pharmacist. You can also report side effects directly via the national reporting system at the address <http://www.aifa.gov.it/content/segnalazioni-reazioni-avverse>. By reporting side effects, you can help provide more information on the safety of this medicine.

SHELF LIFE AND CONSERVATION

Don't use the medicine after the expiry date indicated on the package.

Any special precaution for conservation.

REVISION OF THE LEAFLET BY THE MINISTRY OF HEALTH

December 2019